

Alcohol, Pregnancy and Partner Support

Information for Service Providers

Many women are more likely to avoid alcohol during pregnancy if their partners support them in doing so. Conversely, women find it more difficult to reduce their drinking when their partners oppose that decision or take steps to undermine their efforts. If a woman's partner is supportive of her decision regarding reducing alcohol use during pregnancy, service providers can involve partners in supporting healthy alcohol-free pregnancies.

5 Ways Partners Can Support Women

Service providers can talk to women about what they would find supportive. Some of the things women might find helpful include having their partner:

1 Take a pause from alcohol during the pregnancy by quitting or cutting down. Avoid drinking around her and in social situations when they are together.

2 Tell family, friends and anyone offering her alcohol that there is no known safe time to drink alcohol during pregnancy and no known safe amount.

3 Offer her non-alcoholic drinks (e.g., mocktails) and bring non-alcoholic drinks to social events.

4 Take part in social and recreational activities with her that don't involve alcohol.

5 Help reduce the stress in her life. (You can ask her for specific details of how you might do this.)

Many partners want to support their pregnant partners. Service providers can involve partners, directly and indirectly, by:

- Including information about how partners can support women in both preconception and prenatal materials
- Asking women if they would like to invite their partner or a close family member or friend to their next appointment to discuss how they can support them in achieving an alcohol-free pregnancy, as well as making healthy choices overall
- Sharing suggestions of how soon-to-be parents can be supportive of their pregnant partners. Encourage partners to ask pregnant women what kind of support they need to avoid alcohol, reduce stress and still have fun.

Remember: a partner can be male or female, and some women may not have the support of a partner and may rely on family or friends.

Many partners are interested in thinking about or changing their own alcohol and drug use as they prepare for parenthood. Service providers can provide information and connect partners with community resources. (It's best to do this separately from a woman's care and treatment.)

Created by and reproduced with kind permission from the British Columbia Centre of Excellence for Women's Health

References

1. van der Wulp, N.Y., C. Hoving, and H. de Vries, A qualitative investigation of alcohol use advice during pregnancy: Experiences of Dutch midwives, pregnant women and their partners. *Midwifery*, 2013. In Press.
2. Gearing, R.E., T. McNeill, and F.A. Lozier, Father involvement and fetal alcohol spectrum disorder: Developing best practices. *Fetal Alcohol Research* 2005. 3: p. e14.
3. Mellingen, S., T. Torsheim, and F. Thuen, Changes in alcohol use and relationship satisfaction in Norwegian couples during pregnancy. *Substance Abuse Treatment, Prevention & Policy*, 2013. 8(1): p. 1-11.
4. Everett, K.D., et al, Men's tobacco and alcohol use during and after pregnancy. *American Journal of Men's Health*, 2007. 1(4): p. 317-325.
5. Chang, G., et al, Brief Intervention for Prenatal Alcohol Use: A Randomized Trial. *Obstet Gynecol*, 2005. 105(5 Pt1): p. 991-998.



The Prevention Conversation

It's time to talk about alcohol and pregnancy

