

Common Messages

Guidelines for talking and writing about FASD

2023

Guidelines for Writing and Talking about FASD

The purpose of this document is to assist those writing and talking about FASD – and the issues related to the disability – to use the same language, statistics, and framing of topics. The intended outcome, over time, will be an improved understanding by the reader/listener with consistent and respectful FASD messaging.

A Note About our Process:

This guideline is a “living document” that was collaboratively developed and is updated each year by Canada FASD Research Network researchers, staff, and partners. Annually, we review emerging research in the field to identify new studies that pertain to language use and/or narratives around FASD. When additions are made to the document, we consult with subject matter experts to ensure that our recommendations reflect the most current literature and best practice. We also maintain an ongoing list of topics that arise in conversations between CanFASD staff, researchers, Adult FASD Expert Collaboration Team, Family Advisory Committee, our larger CanFASD member community, and partners to capture evolving messaging in the broader FASD community, particularly related to those with lived experience.

It is important to remember that a “one-size-fits-all” approach to language and messaging may not apply when writing and talking about FASD. The suggestions here are intended to be inclusive and respectful for all individuals with FASD and their families. However, recognizing the diversity and uniqueness of each person with FASD, we always recommend consulting at an individual level to ensure that language choices are reflective and respectful of each person’s experience.

Key Communication Themes:

The following are some overarching recommendations for all messaging about FASD:

- Respect, dignity, and inherent human worth should be promoted among individuals with FASD, women who use alcohol during pregnancy, and their families:
 - FASD awareness or prevention programs and initiatives should avoid messaging that places blame, shame, or guilt.
 - When using imagery, refrain from using pictures of fetuses, pregnant bellies (separated from the person’s body), and naked pregnant people.
 - Fatalistic or deficits-focused terminology should be replaced with person-first, strength-based, and hope-focused language (see the definition below).
 - These [language guides](#) provide additional context and information about dignity promotion for individuals with FASD.
- Always rely on information that is accurate and based on rigorous, high-quality research (i.e., evidence-based).
- Refrain from stating that FASD is “100% preventable” as this statement greatly oversimplifies the issue and can stigmatize individuals and families.

In order to promote a common language about FASD and to minimize misinterpretation of key issues, CanFASD has developed a [standard definition of FASD](#). Standard definitions are needed to ensure consistency in administrative, clinical, and research operations.

The Canada FASD Research Network recommends using the following definition when writing FASD policy, developing supports, and providing services for individuals with FASD across Canada to promote accurate and consistent language when referring to FASD:

Fetal Alcohol Spectrum Disorder (FASD) is a diagnostic term used to describe impacts on the brain and body of individuals prenatally exposed to alcohol.

FASD is a lifelong disability. Individuals with FASD will experience some degree of challenges in their daily living, and need support with motor skills, physical health, learning, memory, attention, communication, emotional regulation, and social skills to reach their full potential.

Each individual with FASD is unique and has areas of both strengths and challenges.

Le Trouble du spectre de l'alcoolisation fœtale (TSAF) est un terme diagnostique utilisé pour décrire les impacts au cerveau et le corps des individus exposés à l'alcool avant la naissance.

Le TSAF est un handicap permanent à vie. Les individus atteints du TSAF connaîtront un certain degré de défis dans leur vie quotidienne et auront besoin d'être supportés dans les domaines suivants: habiletés motrices, santé physique, apprentissage, mémoire, attention, communication, régulation émotionnelle ainsi que les habiletés sociales pour atteindre leur plein potentiel.

Chaque individu atteint du TSAF est unique et présente à la fois des forces et des défis.

PREVALENCE & STATISTICS

Question/Issue	Research Tells Us	Say This	Why This Matters
<p>How many people have FASD?</p>	<p>In Canada</p> <ul style="list-style-type: none"> - In the Greater Toronto Area, researchers estimated the prevalence to be 2-3% among students aged 7-9 years¹ - In Alberta, estimates are 1-4% in the general population² - Teacher-reported prevalence of children <i>diagnosed</i> with FASD in kindergarten ranged from .01 to .22%^{3,4} <p>In the United States</p> <ul style="list-style-type: none"> - Researchers⁵⁻¹¹ have conducted a series of studies with school-aged children in the US, suggesting a conservative estimate of up to 8% in some regions <p>FASD is often under-recognized and identified, so it is likely that prevalence estimates are conservative.</p> <p>For more information:</p> <ul style="list-style-type: none"> - The Prevalence of FASD¹² 	<p>Researchers currently estimate that at least 4% of individuals in Canada have FASD</p> <p>This translates to more than 1.5 million people</p>	<p style="background-color: #800000; color: white; padding: 5px;">Research that helps us better understand how many people might have FASD in Canada, as well as in special populations, is important, in order to identify the scope of the needs and types of services and supports required to assist individuals and families with FASD across the country.</p> <p style="background-color: #003366; color: white; text-align: center; padding: 5px;">Limitations</p> <p>Limitations in prevalence research make it difficult to know the true rate of FASD in Canada. For example, there may be discrepancies in reporting diagnoses, prenatal alcohol exposure may be under-reported due to stigma and fear of judgment, and individual difficulties may not emerge until school years or later. Additional limitations include small samples, unique settings, use of different methodologies, voluntary participation, and estimating the numbers using pre-existing data. Most prevalence studies have been done with children. Estimates vary widely depending on the study approach and population.</p>
<p>Rates of FASD in certain populations</p>	<p>Children in care^{13,14}</p> <ul style="list-style-type: none"> - <i>Canada</i>: up to 11% - <i>International</i>: more than 20% <p>Justice populations¹⁵⁻¹⁸</p> <ul style="list-style-type: none"> - <i>Canada</i>: 10-23% in youth and up to 18% in adult settings - <i>International</i>: 36% in Australia <p>For more information:</p> <ul style="list-style-type: none"> - FASD Prevalence in Special Populations¹⁹ 	<p>Rates of FASD are higher in certain populations, including those involved in the child welfare and justice systems</p>	<p style="background-color: #800000; color: white; padding: 5px;">Service providers in these professions should be knowledgeable about FASD. Screening tools may be especially useful in settings where there are high rates of FASD.</p> <p style="background-color: #003366; color: white; text-align: center; padding: 5px;">Limitations</p> <p>Research in this area is limited and it is suspected that there are higher rates of undiagnosed FASD in other populations.</p>

<p><i>How much alcohol is “too much” during pregnancy?</i></p>	<ul style="list-style-type: none"> - Despite extensive research, there has been no established safe level of alcohol to consume during pregnancy - Alcohol is a neurotoxic and teratogenic substance, which means that it influences fetal development when consumed during pregnancy - Even “low” levels of prenatal alcohol exposure have been shown to have negative effects on long-term development²⁰ - Researchers and clinical experts in maternal and fetal health recommend that no alcohol is best <p>Access this resource for more information:</p> <ul style="list-style-type: none"> - Why do some women drink alcohol during pregnancy? 	<p>Let’s reframe the question to better reflect the message we want to convey:</p> <p><i>“What do we know about alcohol and pregnancy?”</i></p> <p><i>Experts agree that there is no safe level of drinking during pregnancy</i></p> <p><i>It is safest not to drink during pregnancy</i></p>	<p>There are conflicting messages in the media about how much alcohol can be safely consumed during pregnancy, and these messages perpetuate controversy and confusion. People of child-bearing age who are consuming alcohol should use birth control to prevent an unplanned, alcohol exposed pregnancy.</p> <p style="text-align: center;">Limitations</p> <ul style="list-style-type: none"> - Terms like ‘low’ and ‘moderate’ are unclear and subjective - Many people drink before they know they are pregnant, thus it is important to encourage risk reduction with clear and accurate messaging - There are many reasons why women and gender diverse people may drink alcohol during pregnancy. As such, it is important to use non-judgmental messages (e.g., what is “safest”) to convey that there is always opportunity to make a change to reduce risk
<p><i>FASD, mental health, and substance use</i></p>	<ul style="list-style-type: none"> - An estimated 90% of people with FASD experience an additional mental health diagnosis^{21–23} - Some of the most common co-occurring mental health and neurodevelopmental diagnoses among people with FASD are ADHD, intellectual disability, learning disorder, oppositional defiant disorder, conduct disorder, depressive and anxiety disorders, substance use, and risk of suicide²⁴ - Substance use has been reported in ~22-80% of adolescents/adults with FASD²⁵ - An estimated 26% of people with FASD experience past or present suicidal thoughts and behaviours; substance use, trauma, and depression/anxiety are associated with a higher risk of suicidality²⁶ - There is a significant gap in the research on mental health and substance use interventions for people with FASD²⁷ 	<p><i>When unsupported, people with FASD may be more likely to experience high rates of mental health and substance use difficulties</i></p>	<p>Given the high rates of mental health and substance use issues reported among people with FASD, targeted interventions should be designed in this area.</p> <p style="text-align: center;">Limitations</p> <p>More research is needed to better understand and address challenges with mental health/substance use for individuals with FASD across the lifespan. In addition, research designed to understand people with FASD who <i>do not</i> have mental health challenges can provide information about what supports, or protective factors may be important.</p>

<p style="text-align: center;"><i>Social and economic investment</i></p>	<p>FASD has been associated with high social and economic costs:</p> <ul style="list-style-type: none"> - In Alberta:²⁸ <ul style="list-style-type: none"> ○ Long-term economic cost of \$130-400 million/year (predicted number of children born each year) ○ Short-term economic cost of \$48-143 million/year (individuals with FASD) - In Canada:²⁹⁻³¹ <ul style="list-style-type: none"> ○ Cost of supporting those aged 0 to 53 with FASD is \$5.3 billion/year ○ \$1.8 billion for health care, corrections, other direct/indirect costs ○ Total cost \$9.7 billion/year 	<p style="text-align: center;"><i>Replace terms such as “cost” and “burden” with: “investment”</i></p> <p style="text-align: center;"><i>Significant social and economic investment is needed to support adequate resources and services to address prevention, intervention, and diagnosis; and foster better outcomes for people with FASD and their families</i></p>	<p>There is conflicting evidence about true total financial investment associated with FASD, and caution should be taken against framing the disability as a burden.</p>
	<p style="text-align: center;">Limitations</p> <ul style="list-style-type: none"> - Emphasis on the financial investment associated with FASD can perpetuate the stigma experienced by individuals with FASD and their families. - Moreover, previous “cost” estimates do not consider social and cultural investment. - Many studies reflect different financial impacts in society depending on what systems they measure (e.g., medical, education, social services, corrections, prevention/research, productivity loss) - Other social impacts are not measurable (e.g., productivity losses for parents, stress/guilt of mothers, etc.) - We do not know how social and economic investment may be impacted when individuals with FASD and their families receive the services and supports they need - Very little research has been done outside of Canada 		

TERMINOLOGY & LANGUAGE

Question/Issue	Research and Advocacy Efforts Tells Us	Take-Away Message	Why This Matters
<p style="text-align: center;">“Secondary disabilities”</p>	<p>Adolescents and adults with FASD can experience high rates of difficulties in daily living and adverse outcomes, including:^{32–34}</p> <ul style="list-style-type: none"> - School disruption - Sexually inappropriate behaviours - Mental health issues - Substance use challenges - Challenges with independence - Employment difficulties - Housing problems - Trouble with the law - Psychiatric or correctional institutions 	<p style="text-align: center;">Replace “secondary disabilities” with: “adverse outcomes” “difficulties in daily living” “impacts” “associated difficulties” or “risks”</p>	<p style="text-align: center;">Why This Matters</p> <p>Use of the term “secondary” may insinuate that these difficulties are not as prominent or important as the brain- and body-based deficits associated with FASD. Comments about adverse outcomes in FASD should also be considered alongside success and possibility, so as not to bias or limit expectations around life trajectories.</p> <hr/> <p style="text-align: center;">Limitations</p> <p>Many of the challenges described in the literature are not biologically-driven or “disabilities” per se, and are not specific to FASD alone.</p>
<p style="text-align: center;">“Invisible disability”</p>	<p>FASD has commonly been referred to as an “invisible” disability, because the majority of people with FASD have no <i>physical</i> signs of impairment.</p>	<p style="text-align: center;">Replace “invisible” with “hidden”</p>	<p>The use of the term “invisible” can invalidate and stigmatize people with FASD and their families. The term “hidden” more accurately and respectfully reflects the characteristics and experiences of this population.</p>
<p style="text-align: center;">“Disease,” “disorder,” “disability,” “syndrome,” and “condition” used interchangeably</p>	<p>Canadian diagnosticians have moved away from using the term “syndrome” and now use the term “disorder” to refer to FASD.³⁵</p>	<p style="text-align: center;">Replace “disease” and “syndrome” with “disorder” or “disability”</p>	<p>Adoption of the terms “disorder” or “disability” better reflects FASD as a spectrum and will ensure consistent language and clinical understanding.</p>

<p>Speaking respectfully about FASD within Indigenous¹ populations</p> <p>FASD is often still viewed as only an “Indigenous issue”</p>	<ul style="list-style-type: none"> - FASD affects people of all cultural and ethnic backgrounds - Canadian discourse in the 1980s and 1990s perpetuated the stereotype that FASD was an “<i>Indigenous issue</i>”³⁶ - Indigenous women and communities have been continuously subjected to stigma, stereotyping, and surveillance^{37,38} - The Canadian Truth and Reconciliation Recommendations identified FASD in calls to action #33 and #34 - CanFASD along with other groups are working to develop strategies and commitments to address these recommendations: <ul style="list-style-type: none"> o Developing an Indigenous Approach to FASD Prevention³⁹ o TRC Call to Action 34: A Framework for Action⁴⁰ o CanFASD Commitment to Indigenous Partnership, Reconciliatory Research, and Action⁴¹ 	<p>All populations where alcohol is used are at risk for FASD</p> <p>Replace outdated terminology with the term of ‘Indigenous’</p>	<p>FASD is a highly complex disability, intertwined with the social determinants of health and other factors which can affect people from all economic, ethnic, religious, racial, or societal backgrounds.</p> <p style="text-align: center;">Limitations</p> <p>There is a lack of consistent and high-quality research comparing rates of FASD among diverse populations and groups.</p>
<p>Strength-based language and healthy outcomes</p>	<ul style="list-style-type: none"> - Although people with FASD may experience many challenges, they also possess diverse strengths, abilities, and resilience⁴² <ul style="list-style-type: none"> o Strengths Among Individuals with FASD⁴³ - FASD-informed practice includes FASD awareness, safe relationships, and working from a person-centered and strength-based approach^{43,44} - Individuals with FASD should always be referred to using person-first language 	<p>People with FASD are resilient and have many strengths</p> <p>A balanced approach in working with people with FASD that incorporates supports or adaptations is important</p>	<p>Focusing solely on the challenges associated with FASD continues to send the message that these are the main attributes of individuals with FASD. Highlighting strengths in FASD is more conducive to reducing stigma and supporting healthy outcomes.</p> <p style="text-align: center;">Limitations</p> <p>Most of the literature on FASD is deficits-based, and there is a critical need for more research focused on strengths.</p>

¹In Canada, the term ‘Indigenous peoples’ refers to descendants of the peoples of North America who were present before colonization and encompasses three distinct groups: First Nations, Métis, and Inuit.

<p><i>FASD as a “permanent” disability</i></p>	<ul style="list-style-type: none"> - There is no “cure” for FASD, but early intervention and stable home environments are significant protective factors against later life adversity³² - We have moved away from the medical model (which posits that disabilities should be “corrected”) to recognize that individuals with FASD may develop differently but they still develop - Use of the word “treatment” can imply that the effects of FASD are reversible 	<p><i>There is good evidence that early and appropriate support can improve outcomes for people with FASD</i></p>	<p>FASD is a life-long, whole-body disorder, but early identification and intervention can foster healthier outcomes for people with FASD.</p> <p>Although there is no “cure” for FASD, focusing on the permanence of FASD can undercut the fact that intervention can be important and beneficial.</p>
<p><i>“Living with” or “affected by” FASD</i></p>	<ul style="list-style-type: none"> - FASD is not an entity or a person and is not “lived with” - Similarly, stating that someone is “affected by” FASD may imply affliction, distress, or hopelessness 	<p><i>Replace the terms “living with” and “affected by” with “diagnosed with” or “has FASD”</i></p>	<p>Using terms like “living with” or “affected by” FASD may suggest a struggle or burden and minimize the strengths and successes of individuals with FASD.</p>
<p><i>Lived Experience(s) of FASD</i></p>	<ul style="list-style-type: none"> - When describing lived experiences of FASD the terms “in-home” and “in-body” can be used to distinguish the experiences of people who live with an individual who has FASD such as a partner, a caregiver, family, etc. (i.e., “in-home” lived experience) and those who have FASD (i.e., “in-body” lived experience) - Engaging meaningfully with people with in-body and in-home lived experience in all FASD research, policy, and practice allows for the knowledge and voices of those most impacted to be highlighted.⁴⁵ - Involving the perspectives of people with in-home lived experiences of providing support, advocacy, and mentoring to people with FASD, as well as the experiential wisdom of people with in-body lived experience, will help to keep research real, impactful, meaningful, and practical.⁴⁵ 	<p><i>Consider using the terms “in-body” and “in-home” to specify people with lived experience</i></p>	<p>Using these terms may help to distinguish the unique experiences of people with FASD and those with experiences of caregiving or supporting someone with FASD.</p>

<p><i>“Suspected FASD”</i></p>	<ul style="list-style-type: none"> - For some people, the word “suspected” may connote associations with criminal justice involvement. 	<p><i>Replace the term “suspected FASD” with “possible FASD” or someone who “may have FASD”</i></p>	<p>Using the term “suspected FASD” may add to the misperception that FASD is linked to criminal offending and may further stigmatize individuals with FASD.</p>
<p><i>Referring to challenges</i></p>	<ul style="list-style-type: none"> - Individuals with FASD are often referred to as having “significant brain damage” - Terms like “victim,” “suffering,” “damaged child,” and “injured” can further marginalize people with FASD and perpetuate stigma and blame towards parents and family members⁴⁶ - Using the term like “external brain” can further stigmatize and imply that people with FASD do not have a brain - Stigmatizing terms should be replaced by neutral language 	<p><i>Replace terms like “brain damage” with “brain-based differences” “challenges,” “concerns,” “needs” or “difficulties”</i></p>	<p>A balance must be found between acknowledging that significant challenges exist for individuals with FASD and their families, and framing these challenges as permanent and unmalleable. This balance is needed to convey an accurate understanding of FASD while also reducing marginalization and building hope for healthier outcomes.</p>
<p><i>“Won’t” vs “Can’t” vs “How?”</i></p>	<ul style="list-style-type: none"> - There has been a gradual shift from understanding FASD-related learning and behavioural difficulties as willful (i.e., a person ‘won’t’) to understanding these difficulties as resulting from the brain-based impacts of FASD (i.e., a person ‘can’t’). - However, asserting that people with FASD “can’t” contributes to negative stereotypes and expectations, and may perpetuate feelings of hopelessness, self-fulfilling prophecies, and inadequate support.⁴⁷ - Researchers and advocates encourage a continued evolution of this narrative, replacing “can’t” with “how?” That is, <i>how</i> might service providers and others support success with people with FASD?⁴⁸ 	<p><i>Replace terms like “won’t” and “can’t” with “how?”</i></p>	<p>Building on the evolving narrative of “won’t” to “can’t” to “how?” we can reframe our understanding and approaches to intervention and support to better identify ways of facilitating growth and wellbeing for people with FASD.</p>

<p>Parents, caregivers, and families</p>	<ul style="list-style-type: none"> - Parents, caregivers, and families of individuals with FASD experience multiple layers of stigma and marginalization^{46,47,49,50} - Biological parents of children with FASD face particularly high rates of stigma, and may be further marginalized by the use of harmful language - FASD should not be described as having been “caused by” or “the result of” a mother’s alcohol consumption; rather, describe the challenges simply as being associated with prenatal alcohol exposure <p>For more information:</p> <ul style="list-style-type: none"> - FASD Media Guide - Mothers' Experiences of Stigma 	<p>Replace harmful terms such as “alcoholic mother” with person-first terms such as “birth mother, or parent who use(d) substances during pregnancy”</p>	<p>Stigmatizing language still exists in the FASD literature and serves to further marginalize vulnerable individuals and their families. Person-first, strength-based language should replace outdated and harmful messages about parents, caregivers, and families of those with FASD.</p>
<p>Gender-inclusive language</p>	<ul style="list-style-type: none"> - Not all individuals who give birth to a child with FASD will identify as women. In interactions with those who are pregnant, it is important to ask about how they identify their gender and respectfully refer to them using their pronouns. - To be inclusive, clear, and respectful of how all people wish to be referred to, gender additive terms such as “pregnant women and gender diverse individuals” may be considered. When writing generally about those who are pregnant, gender-neutral language such as “pregnant people” and “birthing parent” may also be considered. - At the same time, it is important for some women to be referred to as women or mothers. And for some people, the gender-neutral terms such as “pregnant person” lacks clarity and may be understood to refer to the birth partner not only the pregnant woman. - In many Indigenous cultures “mother” and “mothering” often have particular importance as roles and thus those terms may be important to use. - To ensure language is inclusive and to avoid “othering,” always consult directly with people with lived experience as to what language is appropriate and respectful for them 	<p>To be inclusive of individuals who have conceived or given birth to a child with FASD and do not identify as women, when writing about FASD and PAE consider the use of terms such as “birthing parent(s)” or “women and gender diverse individuals who have given birth”</p>	<p>Gender-inclusive language is needed to reduce gender bias and promote equality across the gender spectrum for parents of individuals with FASD as well as individuals who have FASD themselves.</p> <p style="text-align: center;">Limitations</p> <p>There is a significant gap in the research on gender identity among individuals with FASD, and among people who have given birth to a child with FASD. It is important that individuals see themselves reflected in the language we use, including in research, and to ask people directly how they want to be referred to and be respectful of their identity.</p>

LEGAL ISSUES

Question/Issue	Research Tells Us	Take-Away Message	Why This Matters
<p style="text-align: center;"><i>Involvement in the justice system</i></p>	<ul style="list-style-type: none"> - There are high rates of FASD in youth and adult justice settings,^{17,18} and many people with FASD report experiencing legal trouble^{23,33} - However, not all individuals with FASD end up in trouble with the law - Media portrayals of FASD in the justice system may perpetuate harmful generalizations about criminality in individuals with FASD⁴⁷ - Individuals with FASD also find themselves involved in civil and family law - Individuals with FASD are also victims of crime <ul style="list-style-type: none"> o FASD and the Criminal Justice System: A Review⁵¹ o Victimization in People with FASD⁵² 	<p style="text-align: center;"><i>The challenges and life events experienced by individuals with FASD may lead to increased risk for a range of adverse outcomes when unsupported, including involvement in the justice system</i></p>	<p style="text-align: center;">Gaps in services can leave individuals with FASD and their families unsupported, which can lead to negative outcomes such as justice involvement. However, not all individuals with FASD are justice-involved and we must consider the potentially damaging consequences of associating FASD with justice-involvement.</p> <p style="text-align: center;">Limitations</p> <p>FASD is not the only disability present in justice populations – intellectual disability and psychiatric diagnoses as also over-represented. More research and data are needed in corrections and legal systems to better understand individuals with FASD who are justice-involved</p>
<p style="text-align: center;"><i>Violent crimes and portrayal of FASD in the media</i></p>	<ul style="list-style-type: none"> - There is no research to suggest that justice-involved individuals with FASD commit more <i>violent</i> crimes than justice-involved individuals without FASD - Several studies suggest that justice-involved individuals with FASD are no more likely to commit violent crimes than justice-involved individuals without FASD, but may be more likely to show a “mixed” pattern of offending, with both violent and non-violent crimes^{53,54} 	<p style="text-align: center;"><i>There is no consistent pattern of criminal behaviour that has been identified for justice-involved individuals with FASD</i></p>	<p style="text-align: center;">Media portrayals of violent crime and FASD tend to be sensationalized and can serve to misrepresent justice-involved individuals with FASD.</p> <p style="text-align: center;">Limitations</p> <p>The media often reports stories in which people with FASD are involved in <i>violent</i> crimes, giving FASD a violent connotation. We need more information about justice-involvement in FASD to tell the whole story.</p>
<p style="text-align: center;"><i>Alternative justice practices</i></p>	<p>There are numerous initiatives across the country to better address FASD in the justice system (e.g., restorative justice, diversion programs, conditional or alternative sentencing, mental health courts) but very little research has been done to examine long term outcomes of such programs.</p>	<p style="text-align: center;"><i>Alternative justice practices may be promising for working with some justice-involved individuals with FASD, but more research is needed</i></p>	<p style="text-align: center;">The traditional justice system is not adequately set up to manage justice-involved individuals with FASD, but more research is needed to assess the long-term impact of alternative justice practices.</p>

EMERGING RESEARCH

Question/Issue	Research Tells Us	Take-Away Message	Why This Matters
<p><i>“If a man drinks, will the sperm cause FASD?”</i></p>	<p>There is very little research on the connection between FASD and alcohol consumption by male partners (or gender diverse partners who can contribute to conceptions) during the preconception or prenatal period (see Preconception paternal ethanol exposures induce alcohol-related craniofacial growth deficiencies in fetal offspring). However, male/gender-diverse partner alcohol consumption has been associated with:⁵⁵</p> <ul style="list-style-type: none"> - Higher levels of maternal alcohol consumption during pregnancy - Lower levels of success in infertility treatments - Lower infant birth weight and gestational age - Higher rates of ventricular malformations and abnormal situs in infants <p>For more information:</p> <ul style="list-style-type: none"> - The Role of Partners in Fetal Alcohol Spectrum Disorder Prevention⁵⁶ - Genetic and Epigenetic Perspectives on the Role of Fathers in Fetal Alcohol Spectrum Disorder⁵⁷ 	<p><i>Fathers’ drinking does not result in FASD, but it has an important influence on mothers’ consumption and the healthy infant and child development</i></p>	<p>Fathers and all partners have an important responsibility in supporting their pregnant partners to stay healthy during pregnancy, and in the long-term health of their children</p> <p style="text-align: center;">Limitations</p> <p>This is a very new area of research, and more studies are needed to understand the role of fathers’ alcohol consumption in healthy child development.</p>
<p><i>How should the role of the DSM-5’s Neurobehavioral Disorder Associated with Prenatal Alcohol Exposure be described in relation to the Canadian Diagnostic Guideline?</i></p>	<ul style="list-style-type: none"> - The DSM-5 defines ND-PAE as a condition characterized by PAE and impairment in neurocognitive, self-regulation, and adaptive functioning⁵⁸ - It is <i>not</i> a formal diagnosis, but proposed as a “condition for further study” - Researchers recommend a comprehensive assessment conducted by a multidisciplinary team, regardless of diagnostic approach^{35,59} <p>For more information:</p> <ul style="list-style-type: none"> - Multidisciplinary Team Training for Diagnosis of FASD: An Online Curriculum 	<p><i>We encourage practitioners and clinicians working with individuals with PAE to receive training on the 2015 Canadian FASD diagnostic guideline</i></p>	<p>In Canada, the 2015 FASD Diagnostic Guideline is primarily used to assess and diagnose individuals with PAE. The Canadian Guideline supports a multidisciplinary approach to diagnosis which is important for a full understanding of the neurocognitive impairments and the required interventions and supports.</p> <p style="text-align: center;">Limitations</p> <p>More research is needed to compare and contrast the different diagnostic systems used across the world.</p>

<p><i>FASD and Autism Spectrum Disorder are often compared</i></p>	<ul style="list-style-type: none"> - FASD and ASD are both neurodevelopmental and spectrum disorders, but their etiologies differ – there is a known cause of FASD, while the cause of ASD is unclear - Despite some overlapping symptoms (e.g., social and communication difficulties), FASD and ASD have distinct presentations in clinical and daily settings⁶⁰ - Families of individuals with FASD/ASD have different experiences⁶¹ 	<p><i>FASD and ASD are both neurodevelopmental spectrum disorders with some overlapping symptoms, but distinct presentations</i></p>	<p>FASD and ASD share some similarities, but they are managed and experienced differently.</p>
		<p>Limitations</p>	
		<p>Some emerging epigenetic research explores the link between FASD and ASD, but little is known about the association between the two disorders.</p>	
<p><i>Life expectancy and hospitalization</i></p>	<ul style="list-style-type: none"> - Some researchers have reported that individuals with PAE and FASD may experience a reduced life expectancy,⁶² accelerated aging processes,⁶³ and increased risk for hospitalization and mortality⁶⁴ - This research is preliminary and should therefore be interpreted with caution 	<p><i>Individuals with FASD may be at risk for poorer medical outcomes, but these are not guaranteed</i></p> <p><i>Access to adequate supports and services can alter long-term trajectories</i></p>	<p>Fatalistic messaging can increase stigma and hopelessness for individuals with FASD. Messages related to life expectancy should be appropriately contextualized and accompanied by messages of healthy outcomes.</p>
		<p>Limitations</p>	
		<p>Research in this area is in its infancy, limited geographically, and lacks generalizability.</p>	

OTHER ISSUES

Question/Issue	Research Tells Us	Take-Away Message	Why This Matters
Developmental age versus chronological age	<ul style="list-style-type: none"> - People with FASD often function at a level younger than their chronological age⁶⁵ - Functioning may also be uneven across domains, with high abilities in some areas but not others⁶⁶ - With increasing age, societal expectations of autonomy and independence may be particularly challenging for those with FASD⁶¹ 	<p><i>Although age descriptors can be helpful in describing an individuals' situation, moving away from this framing will better support a strength-based narrative of FASD</i></p>	<p style="background-color: #800000; color: white; padding: 5px;">Individuals with FASD have strengths as well as challenges that should both be taken into consideration to best understand and meet needs.</p> <p style="background-color: #003366; color: white; text-align: center; padding: 5px;">Limitations</p> <p>Focusing solely on developmental age may lead us to neglect the fact that, with support, individuals with FASD continue to develop and achieve success.</p>
Neurodiversity	<ul style="list-style-type: none"> - At its core, neurodiversity is a concept that represents the wide variety of differences among humanity. - Neurodiversity is a concept encompassing both individual level (i.e., identity) as well as systemic level (i.e., political movement) considerations.⁶⁷ It is a relatively new phenomenon and as such a clear definition and understanding of neurodiversity is not currently shared.⁶⁷ - The concept of neurodiversity does not disregard the uniqueness of individuals with FASD. The terms neurodiversity and FASD are not interchangeable, but each term is valuable. - It is important to respect how individual people wish to be identified and referred to. 	<p><i>Neurodiversity is a concept with increased attention at the individual and systemic levels. The terms FASD and neurodiversity are not interchangeable but both terms are valuable. Adopting a both/and approach to FASD and neurodiversity allows space for understanding of the unique elements of FASD, while also providing space for the valuing of those differences</i></p>	<p style="background-color: #800000; color: white; padding: 5px;">Neurodiversity as a concept for both individual identity and a social movement has received increased attention. Understanding where FASD fits into this framing is an important continued consideration. It is always important to respect how people wish to be identified and referred to.</p> <p style="background-color: #003366; color: white; text-align: center; padding: 5px;">Limitations</p> <p>Additional research is needed to understand how FASD fits within the context of neurodiversity, including engaging with people with in-body and in-home lived experience.</p>
Interdependence	<ul style="list-style-type: none"> - Whereas <i>independence</i> is the idea that we should be able to do everything on our own, <i>interdependence</i> is the notion that we all rely on one another, and there is a spectrum within which we access support to navigate different elements of functioning. - We are all interdependent to some extent and it is important to normalize this truth, and find the appropriate level of 	<p><i>Consider replacing the idea of "independence" with the concept of "interdependence," whereby every individual relies on people</i></p>	<p>Interdependence is a reasonable and achievable goal for all people with FASD. Recognizing this goal while understanding that interdependence will look different for all people is important. Needs arounds interdependence should be considered throughout the lifespan, and maintained even when an individual is living well.</p>

	<p>interdependence on an individual basis, which may differ between people and tasks.⁴⁸</p> <ul style="list-style-type: none"> - Priorities for and levels of interdependence can differ depending on an individual's values, perspectives, abilities, and needs as well as a person's family or cultural values. - It is important that when people with FASD begin to thrive within networks of interdependence, supports not be taken away. 	<p><i>and systems to varying degrees, and that interdependence can mean different things to each person with FASD.</i></p>	<p style="text-align: center;">Limitations</p> <p>Additional research is needed to explore needs and priorities around interdependence and how best to support people with FASD throughout their lifespan.</p>
--	---	---	---

References

1. Popova S, Lange S, Chudley A, et al. *World Health Organization International Study on the Prevalence of Fetal Alcohol Spectrum Disorder (FASD) Canadian Component*. Toronto, ON. Retrieved from <https://canfasd.ca/wp-content/uploads/sites/35/2018/05/2018-Popova-WHO-FASD-Prevalence-Report.pdf>
2. Thanh NX, Jonsson E, Salmon A, et al. Incidence and prevalence of Fetal Alcohol Spectrum Disorder by sex and age group in Alberta, Canada. *J Popul Ther Clin Pharmacol* 2014;21(3):e395-404. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/25381628>
3. Palmeter S, Probert A, Lagacé C. FASD prevalence among children and youth: Results from the 2019 Canadian Health Survey on Children and Youth. *Heal Promot Chronic Dis Prev Canada Res Policy Pract* 2021;41(9):272–276. [doi:10.24095/hpcdp.40.9.05](https://doi.org/10.24095/hpcdp.40.9.05)
4. Pei J, Reid-Westoby C, Siddiqua A, et al. Teacher-reported prevalence of FASD in kindergarten in Canada: Association with child development and problems at home. *J Autism Dev Disord* 2021;51(2):433–443. <https://doi.org/10.1007/s10803-020-04545-w>
5. May PA, Baete A, Russo J, et al. Prevalence and characteristics of Fetal Alcohol Spectrum Disorders. *Pediatrics* 2014;134(5):855–866. [doi:10.1542/peds.2013-3319](https://doi.org/10.1542/peds.2013-3319)
6. May PA, Chambers CD, Kalberg WO, et al. Prevalence of Fetal Alcohol Spectrum Disorders in 4 US communities. *JAMA* 2018;319(5):474–482. [doi:10.1001/jama.2017.21896](https://doi.org/10.1001/jama.2017.21896)
7. May PA, Keaster C, Bozeman R, et al. Prevalence and characteristics of Fetal Alcohol Syndrome and Partial Fetal Alcohol Syndrome in a Rocky Mountain region city. *Drug Alcohol Depend* 2015;1(115):118–127. [doi:10.1016/j.drugalcdep.2015.08.006](https://doi.org/10.1016/j.drugalcdep.2015.08.006)
8. May PA, Hasken JM, Baete A, et al. Fetal Alcohol Spectrum Disorders in a Midwestern City: Child characteristics, maternal risk traits, and prevalence. *Alcohol Clin Exp Res* 2020;44(4):919–938. [doi:10.1111/acer.14314](https://doi.org/10.1111/acer.14314)
9. May PA, Hasken JM, Bozeman R, et al. Fetal Alcohol Spectrum Disorders in a Rocky Mountain Region City: Child characteristics, maternal risk traits, and prevalence. *Alcohol Clin Exp Res* 2020;44(4):900–918. [doi:10.1111/acer.14315](https://doi.org/10.1111/acer.14315)
10. May PA, Hasken JM, Stegall JM, et al. Fetal Alcohol Spectrum Disorders in a Southeastern County of the United States: Child characteristics and maternal risk traits. *Alcohol Clin Exp Res* 2020;44(4):939–959. [doi:10.1111/acer.14313](https://doi.org/10.1111/acer.14313)
11. May PA, Hasken JM, Hooper SR, et al. Estimating the community prevalence, child traits, and maternal risk factors of Fetal Alcohol Spectrum Disorders (FASD) from a random sample of school children. *Drug Alcohol Depend*;227. [doi:10.1016/j.drugalcdep.2021.108918](https://doi.org/10.1016/j.drugalcdep.2021.108918)
12. Flannigan K, Unsworth K, Harding K. *The prevalence of Fetal Alcohol Spectrum Disorder*. 2018. *CanFASD*. Retrieved from: <https://canfasd.ca/wp-content/uploads/publications/Prevalence-1-Issue-Paper-FINAL.pdf>
13. Fuchs D, Burnside L. *A tri-province initiative to expand understanding of costs, services and prevention of a public health issue: Fetal Alcohol Spectrum Disorder and children/youth in care [2010-2014]: Study on the prevalence of FASD in Canadian child welfare settings: Final Report*. Manitoba, MB 2014.
14. Lange S, Shield K, Rehm J, et al. Prevalence of Fetal Alcohol Spectrum Disorders in child care settings: A meta-analysis. *Pediatrics* 2013;132(4):e980–e985. [doi:10.1542/peds.2013-0066](https://doi.org/10.1542/peds.2013-0066)
15. Bower C, Watkins RE, Mutch RC, et al. Fetal Alcohol Spectrum Disorder and youth justice: A prevalence study among young people sentenced to detention in Western Australia. *BMJ Open*;8(2). [doi:10.1097/00004703-199910000-00012](https://doi.org/10.1097/00004703-199910000-00012)
16. Fast DK, Conry J, Looock CA. Identifying Fetal Alcohol Syndrome among youth in the criminal justice system. *J Dev Behav Pediatr* 1999;20(5):370–372.
17. McLachlan K, McNeil A, Pei J, et al. Prevalence and characteristics of adults with Fetal Alcohol Spectrum Disorder in corrections: A Canadian case ascertainment study. *BMC Public Health*;19(1). [doi:10.1186/s12889-018-6292-x](https://doi.org/10.1186/s12889-018-6292-x)
18. Popova S, Lange S, Bekmuradov D, et al. Fetal Alcohol Spectrum Disorder prevalence estimates in correctional systems: A systematic literature review. *Can J Public Heal* 2011;102(5):336–340. [doi:10.1007/BF03404172](https://doi.org/10.1007/BF03404172)
19. Flannigan K, Unsworth K, Harding K. *FASD prevalence in special populations*. 2018. *CanFASD*. Retrieved from: <https://canfasd.ca/wp-content/uploads/publications/Prevalence-2-Issue-Paper-FINAL.pdf>
20. Day NL, Helsel A, Sonon K, et al. The association between prenatal alcohol exposure and behavior at 22 years of age. *Alcohol Clin Exp Res* 2013;37(7):1171–1178. [doi:10.1111/acer.12073](https://doi.org/10.1111/acer.12073)
21. O’Connor M, Paley B. Psychiatric conditions associated with prenatal alcohol exposure. *Dev Disabil Res Rev* 2009;15(3):225–234. [doi:10.1002/ddrr.74](https://doi.org/10.1002/ddrr.74)
22. Pei J, Denys K, Hughes J, et al. Mental health issues in Fetal Alcohol Spectrum Disorder. *J Ment Heal* 2011;20(5):473–483. [doi:10.3109/09638237.2011.577113](https://doi.org/10.3109/09638237.2011.577113)
23. Streissguth A, Barr H, Kogan J, et al. *Understanding the occurrence of secondary disabilities in clients with Fetal Alcohol Syndrome (FAS) and Fetal Alcohol Effects (FAE)*. Seattle, WA 1996. Retrieved from: <http://lib.adai.uw.edu/pubs/bk2698.pdf>
24. Weyrauch D, Schwartz M, Hart B, et al. Comorbid mental disorders in Fetal Alcohol Spectrum Disorders: A systematic review. *J Dev Behav Pediatr* 2017;38(4):283–291. [doi:10.1097/DBP.0000000000000440](https://doi.org/10.1097/DBP.0000000000000440)

25. Popova S, Lange S, Burd L, et al. Cost of specialized addiction treatment of clients with Fetal Alcohol Spectrum Disorder in Canada. *BMC Public Health* 2013;13(1):1–11. [doi:10.1186/1471-2458-13-570](https://doi.org/10.1186/1471-2458-13-570)
26. Flannigan K, McMorris C, Ewasiuk A, et al. Suicidality and associated factors among individuals assessed for Fetal Alcohol Spectrum Disorder across the lifespan in Canada. *Can J Psych* 2022;67(5):361-370. [doi:10.1177/07067437211053288](https://doi.org/10.1177/07067437211053288)
27. Flannigan K, Coons-Harding KD, Anderson T, et al. A systematic review of interventions to improve mental health and substance use outcomes for individuals with prenatal alcohol exposure and Fetal Alcohol Spectrum Disorder. *Alcohol Clin Exp Res* 2020;44(12):2401–2430. [doi:10.1111/acer.14490](https://doi.org/10.1111/acer.14490)
28. Thanh NX, Jonsson E. Costs of Fetal Alcohol Spectrum Disorder in Alberta, Canada. *Can J Clin Pharmacol* 2009;16(1):e80–e90. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/19151424>
29. Stade B, Ali A, Bennet D, et al. The burden of prenatal alcohol exposure: Revised measurement of cost. *Can J Clin Pharmacol* 2016;16(1):e91–e102. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/19168935>
30. Popova S, Lange S, Burd L, et al. Cost attributable to Fetal Alcohol Spectrum Disorder in the Canadian correctional system. *Int J Law Psychiatry* 2015;41:76–81. [doi:10.1016/j.ijlp.2015.03.010](https://doi.org/10.1016/j.ijlp.2015.03.010)
31. Thanh NX, Jonsson E. Costs of Fetal Alcohol Spectrum Disorder in the criminal justice system. *J Popul Ther Clin Pharmacol* 2015;22(1):e125–e131. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/26072470>
32. Streissguth AP, Bookstein FL, Barr HM, et al. Risk factors for adverse life outcomes in Fetal Alcohol Syndrome and Fetal Alcohol Effects. *J Dev Behav Pediatr* 2004;25(4):228–238. [doi:10.1097/00004703-200408000-00002](https://doi.org/10.1097/00004703-200408000-00002)
33. McLachlan K, Flannigan K, Temple V, et al. Difficulties in daily living experiences by adolescents, transition-aged youth, and adults with Fetal Alcohol Spectrum Disorder. *Alcohol Clin Exp Res* 2020;44(8):1609–1624. [doi:10.1111/acer.14385](https://doi.org/10.1111/acer.14385)
34. Rangmar J, Hjern A, Vinnerljung B, et al. Psychosocial outcomes of Fetal Alcohol Syndrome in adulthood. *Pediatrics* 2015;135(1):e52–e58. [doi:10.1542/peds.2014-1915](https://doi.org/10.1542/peds.2014-1915)
35. Cook JL, Green CR, Lilley CM, et al. Fetal Alcohol Spectrum Disorder: A guideline for diagnosis across the lifespan. *Can Med Assoc J* 2015;188(3):191–197. [doi:10.1503/cmaj.141593](https://doi.org/10.1503/cmaj.141593)
36. McKenzie HA, Dell CA, Fornssler B. Understanding addictions among Indigenous people through social determinants of health frameworks and strength-based approaches: A review of the research literature from 2013 to 2016. *Curr Addict Reports* 2016;3:378–386. [doi:10.1007/s40429-016-0116-9](https://doi.org/10.1007/s40429-016-0116-9)
37. Tait C. “Tip of the iceberg”: The “making” of Fetal Alcohol Syndrome in Canada. Doctoral Thesis, McGill University; 2003. Retrieved from <http://www.namhr.ca/media/docs/lega4eb233149a533-tait-2003.pdf>
38. Yousefi N, Chaufan C. “Think before you drink”: Challenging narratives on Foetal Alcohol Spectrum Disorder and Indigeneity in Canada. *Health (London)* 2021; [doi:10.1177/13634593211038527](https://doi.org/10.1177/13634593211038527)
39. CanFASD, Thunderbird Partnership Foundation, Centre of Excellence for Women's Health. *Consensus statement: Eight tenets for enacting the Truth and Reconciliation Commission's Call to Action #33*. 2017. Retrieved from http://bcccewh.bc.ca/wp-content/uploads/2017/09/Indigenous_approach_FASD.pdf
40. Stewart M, Glowatski K. *Truth & Conciliation Call to Action #34: A framework for action*. CanFASD. Retrieved from <https://canfasd.ca/wp-content/uploads/sites/35/2018/10/TRC34-A-Framework-for-Action.pdf>
41. CanFASD. *Canada FASD Research Network's commitment to Indigenous partnership, reconciliatory research, and action*. Retrieved from <https://canfasd.ca/wp-content/uploads/publications/CanFASDs-Commitment-to-Indigenous-Partnership-Reconciliatory-Research-and-Action.pdf>
42. Olson HC, Sparrow J. A shift in perspective on secondary disabilities in Fetal Alcohol Spectrum Disorders. *Alcohol Clin Exp Res* 2021;45(5):916–921. [doi:10.1111/acer.14586](https://doi.org/10.1111/acer.14586)
43. Flannigan K, Wrath A, Ritter C, et al. Balancing the story of Fetal Alcohol Spectrum Disorder: A narrative review of the literature on strengths. *Alcohol Clin Exp Res* 2021;45(12):2448–2464. [doi:10.1111/acer.14733](https://doi.org/10.1111/acer.14733)
44. Rutman D. Becoming FASD informed: Strengthening practice and programs working with women with FASD. *Subst Abus Res Treat* 2016;10 (Suppl1):13–20. [doi:10.4137/SART.S34543](https://doi.org/10.4137/SART.S34543)
45. Reid D, Beland W, Richardson L, et al. What if? Incorporating the voices of those with lived experience to change the focus of Fetal Alcohol Spectrum Disorder research. *J Fetal Alcohol Spectr Disord* 2022;4(SP1):e162–e170. [doi:10.22374/jfasd.v4iSP1.16](https://doi.org/10.22374/jfasd.v4iSP1.16)
46. Roozen S, Stutterheim SE, Bos AER, et al. Understanding the social stigma of Fetal Alcohol Spectrum Disorders: From theory to interventions. *Found Sci* 2020;27:753–771. [doi:10.1007/s10699-020-09676-y](https://doi.org/10.1007/s10699-020-09676-y)
47. Aspler J, Zizzo N, Bell E, et al. Stigmatisation, exaggeration, and contradiction: An analysis of scientific and clinical content in Canadian print media discourse about Fetal Alcohol Spectrum Disorder. *Can J Bioeth* 2019;2(2):23–35. [doi:10.7202/1058140ar](https://doi.org/10.7202/1058140ar)
48. Pei J, Kapasi A, Kennedy KE, et al. Towards Health Outcomes for Individuals with FASD. CanFASD 2019. Retrieved from: <https://canfasd.ca/wp-content/uploads/publications/Final-Towards-Healthy-Outcomes-Document-with-links.pdf>
49. Bell E, Andrew G, Di Pietro N, et al. It's a shame! Stigma against Fetal Alcohol Spectrum Disorder: Examining the ethical implications for public health practices and policies. *Public Health Ethics* 2016;9(1):65–77. [doi:10.1093/phe/phv012](https://doi.org/10.1093/phe/phv012)
50. Corrigan PW, Lara JL, Shah BB, et al. The public stigma of birth mothers of children with Fetal Alcohol Spectrum Disorder. *Alcohol Clin Exp Res* 2017;41(6):1166–1173. [doi:10.1111/acer.13381](https://doi.org/10.1111/acer.13381)
51. Pei J, Flannigan K, Keller S, et al. Fetal Alcohol Spectrum Disorder and the criminal justice system: A review. *J Ment Heal Clin Psychol* 2018;2(4):48–52. Retrieved from: <https://canfasd.ca/wp-content/uploads/sites/35/2018/08/FASD-Justice-Review-Issue-Paper-FINAL.pdf>

52. Glowatski K, Stewart M. *Victimization in people with FASD*. CanFASD. Retrieved from: <https://canfasd.ca/wp-content/uploads/2018/07/FASD-and-Victimization-FINAL.pdf>
53. McLachlan K. An examination of the abilities, risks, and needs of adolescents and young adults with Fetal Alcohol Spectrum Disorder (FASD) in the criminal justice system. Doctoral Thesis, Simon Fraser University; 2012. Retrieved from: <https://summit.sfu.ca/item/12102>
54. MacPherson PH, Chudley AE, Grant BA. Fetal Alcohol Spectrum Disorder in a correctional population: Prevalence, screening and characteristics, Research Report R-247. Correctional Service of Canada. Retrieved from <http://www.csc-scc.gc.ca/research/005008-0247-eng.shtml>
55. McBride N, Johnson S. Fathers' role in alcohol-exposed pregnancies: Systematic review of human studies. *Am J Prev Med* 2016;51(2):240–248. [doi:10.1016/j.amepre.2016.02.009](https://doi.org/10.1016/j.amepre.2016.02.009)
56. Wolfson L, Harding K, Poole N. *The role of partners in Fetal Alcohol Spectrum Disorder prevention*. CanFASD 2019. Retrieved from <https://canfasd.ca/wp-content/uploads/publications/the-role-of-partners-in-fetal-alcohol-spectrum-disorder-prevention.pdf>
57. Liyanage-Zachariah V, Harding K. *Genetic and epigenetic perspectives on the role of fathers in Fetal Alcohol Spectrum Disorder*. CanFASD 2019. Retrieved from: <https://canfasd.ca/wp-content/uploads/publications/Fathers-Role-1-Issue-Paper-Final.pdf>
58. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association; 2013.
59. Doyle LR, Mattson SM. Neurobehavioral Disorder Associated with Prenatal Alcohol Exposure (ND-PAE): Review of evidence and guidelines for assessment. *Curr Dev Disord Reports* 2015;2(3):175–186. [doi:10.1007/s40474-015-0054-6](https://doi.org/10.1007/s40474-015-0054-6)
60. Temple V, Cook JL, Unsworth K, et al. Prenatal alcohol exposure and Autism Spectrum Disorder in 39 children and adults: Examination of behavioural and cognitive profiles. *J Ment Health Res Intellect Disabil* 2021;14(1):107–121. [doi:10.1080/19315864.2020.1864688](https://doi.org/10.1080/19315864.2020.1864688)
61. Watson S, Hayes S, Radford-Paz E, et al. “I’m hoping, I’m hoping...” Thoughts about the future from families of children with autism or fetal alcohol spectrum disorder in Ontario. *J Dev Disabil* 2013;19(3):76–93.
62. Thanh NX, Jonsson E. Life Expectancy of People with Fetal Alcohol Syndrome. *J Popul Ther Clin Pharmacol* 2016;23(1):e53–e59. Retrieved from <http://psycnet.apa.org/record/2014-06811-007>
63. Okazaki S, Otsuka I, Shinko Y, et al. Epigenetic clock analysis in children with Fetal Alcohol Spectrum Disorder. *Alcohol Clin Exp Res* 2021;45(2):329–337. [doi:10.1111/acer.14532](https://doi.org/10.1111/acer.14532)
64. Oh SS, Kim YJ, S J, et al. Hospitalizations and mortality among patients with Fetal Alcohol Spectrum Disorders: A prospective study. *Sci Rep*;10(1). [doi:10.1038/s41598-020-76406-6](https://doi.org/10.1038/s41598-020-76406-6)
65. Bennett B. *Life stages and transitions*. Paper presented at the Fetal Alcohol Spectrum Disorder (FASD): Across the lifespan: Proceedings from an IHE consensus development conference; 2009.
66. MOFAS. *Raising children with Fetal Alcohol Spectrum Disorders (FASD)*. 2017. Retrieved from: <https://edmontonfetalalcoholnetwork.org/2017/05/29/raising-children-with-fetal-alcohol-spectrum-disorders-fasd/>
67. Harding K, Pei J, Richardson L. *Neurodiversity and FASD*. CanFASD 2023. Retrieved from: <https://canfasd.ca/wp-content/uploads/publications/Neurodiversity-and-FASD.pdf>



Thank you to all of the CanFASD researchers and staff who provided valuable input into this document.

If you have comments or suggestions, please send them to Audrey.McFarlane@canfasd.ca